


MALDI/TOF MS for Chemical Analysis of Fingerprint Residues using **Conventional Fingerprint Development Methods**

OVERVIEW

- from a fingerprint using conventional methods.



ACKNOWLEDGEMENTS

This research is supported in part by the FBI's Visiting Scientist Program, an educational opportunity administered by the Oak Ridge Institute for Science and Education. Names of commercial manufacturers are provided for identification purposes only, and inclusion does not imply endorsement of the manufacturer, or its products or services by the FBI. The views expressed are those of the authors' and do not necessarily reflect the official policy or position of the FBI or the U.S. Government. Benton M, et al. (2009) Surf. Interface Anal.:42:378-385 Rowell F et al (2009) Analyst 4:701-707

Kimberly Kaplan^{*1,2}, Mark L Miller², and Marc A LeBeau³

fingerprints


¹Oak Ridge Institute for Science and Education, Oak Ridge, Tennessee, ²Counterterrorism and Forensic Science Research Unit, ³Scientific Analysis Section, Federal Bureau of Investigation, Quantico, Virginia USA, Laboratory Division, Federal Bureau of Investigation, Quantico, Virginia USA



Figure 4: MALDI/TOF MS images of control and ibuprofen fingerprints developed with four different powders. Images for [M-H]⁻ shows chemical interference in the controls and ibuprofen handled latent

RESULTS

POWDER COMPARISON



reduced after lifting.

cyanoacrylate fuming followed by application of four different powders. Compared to Figure 4 chemical interference is reduced with cyanoacrylate fuming and powdering



